

# Probabilistic Teleportation of Arbitrary Three-Particle GHZ and W State

WANG Haozhen, TAN Xiaoqing\*, XU Qingshan, and BAO Daipengwei  
(College of Information Science and Technology, Jinan University, Guangzhou 510632, China)

**Abstract** A new quantum channel composed of two multi-particle partially entangled states for quantum teleportation (QT) is constructed. Two explicit controlled probabilistic teleportation tripartite schemes are proposed for arbitrary three-particle partially entangled GHZ state and W state. The arbitrary GHZ state and W state can be teleported with a certain probability to the receiver who introduces an auxiliary particle and performs the corresponding unitary operation, with the help of the measurement of Bell-state and single-particle from the sender and the controller. The overall success probabilities of two QT schemes are both  $8d^2n^2$  ( $d$  and  $n$  are the coefficients of the quantum channel).

**Key words** Bell state measurement; partially entangled state; quantum teleportation; unitary operation

## 任意三粒子 GHZ 态和 W 态的概率隐形传态



王浩圳, 谭晓青\*, 许青山, 包戴鹏伟

(暨南大学 信息科学技术学院, 广州 510632)

**摘要** 构造了一个全新的由两个多粒子部分纠缠态组成的用于量子隐形传态 (QT) 的量子信道。利用构造的量子信道, 提出了两个任意三粒子 GHZ 态和 W 态的受控概率隐形传态的三方方案。发送者执行贝尔态测量和单粒子测量, 控制者执行贝尔态测量, 接收者引入一个辅助粒子并执行相应酉操作, 可实现任意三粒子 GHZ 态和 W 态的概率隐形传态。两个 QT 方案的总成功概率都是  $8d^2n^2$  (其中  $d$  和  $n$  是量子信道的系数)。

**关键词** 贝尔态测量; 部分纠缠态; 量子隐形传态; 酉操作

中图分类号 TN918.1; O431.2

文献标志码 A

DOI 10.12178/1001-0548.2022356

Quantum entanglement is widely used in quantum communication, such as quantum key distribution<sup>[1-3]</sup>, quantum superdense coding<sup>[4]</sup>, quantum secure direct communication<sup>[5-7]</sup>, and quantum teleportation<sup>[8-12]</sup>. The transmission of quantum state could be implemented by quantum teleportation (QT) with the large information capacity and high reliability, but could not be accomplished by pure complete classical methods or pure quantum methods.

In 1993, Ref. [8] firstly presented a complete QT scheme to transfer an arbitrary unknown single-particle to a distant receiver from a sender by using maximally entangled Bell states. Ref. [13] extended the discrete variable teleportation theory of Ref. [8] and conceptually defined the teleportation of continuous variables in 1994. Ref. [14] successfully achieved the single-particle quantum teleportation experimentally in 1997. In 1998, Ref. [15], based on

Received date: 2022-10-20; Revised date: 2023-02-23

收稿日期: 2022-10-20; 修回日期: 2023-02-23

Foundation item: National Natural Science Foundation of China (62032009); Natural Science Foundation of Guangdong Province of China (2021A1515011440); Major Program of Guangdong Basic and Applied Research (2019B030302008)

基金项目: 国家自然科学基金重点项目 (62032009); 广东省自然科学基金 (2021A1515011440); 广东省基础与应用基础研究重大项目 (2019B030302008)

Biography: WANG Haozhen, his research interests include quantum communication.

作者简介: 王浩圳, 主要从事量子通信方面的研究。

\*Corresponding author's E-mail: ttanxq@jnu.edu.cn

\*通信作者 E-mail: ttanxq@jnu.edu.cn

Ref. [11], realized the teleportation of the continuous coherent light field by using the two-mode squeezed vacuum field. With the development of technology, the transmitted particles have expanded from the initial single-particle state<sup>[16-17]</sup> to the multi-particle state<sup>[18-19]</sup>, and the teleportation channels have also expanded from the initial two-particle state to the multi-particle state<sup>[20]</sup>, which composed of the W state, GHZ state, or some other combination of states.

In a practical situation, the influence of the environment and the inevitable occurrence of decoherence make it impossible for entangled channels to be in an ideal, noiseless, and maximally entangled state. Evenmore, they may be in non-maximally entangled states or even mixed states. If these quantum channels are not in maximally entangled state, it is impossible to transmit particles with perfect probability and fidelity. However, it has been shown that QT with the perfect fidelity can also be achieved by using non-maximally entangled states but with a probability less than 1, so it turns out to be possible to use the non-maximally entangled basis as the measurement basis. This new idea of QT is called probabilistic quantum teleportation (PQT)<sup>[21-22]</sup>.

Subsequently, QT has been further developed, and the QT scheme has also been generalized to the controlled quantum teleportation (CQT)<sup>[23]</sup> and  $N$  particles QT<sup>[24]</sup> by using various kinds of quantum channels and the design of the process. There are many CQT schemes<sup>[25-26]</sup>, such as some special bidirectional controllable quantum teleportation (BCQT)<sup>[27-29]</sup> and cyclic controllable quantum teleportation (CCQT)<sup>[30-32]</sup>.

We propose two controllable probabilistic teleportation schemes for an arbitrary three-particle GHZ state or W state in this paper, by using a partially entangled four-particle state and a partially entangled three-particle state as the QT channel. At the same time, our schemes can implement teleportation from a sender to either one of the two receivers. To recover the unknown state, apart from the sender, the only requirement for the controller is to perform Bell-state measurement, and the receiver merely needs to have the ability to perform the unitary operation with the help of the controller. The overall success probabilities

of our two schemes are both  $8d^2n^2$ . The probability of success can attain 100% if the channel is in its maximally entangled state.

## 1 Probabilistic Teleportation of Arbitrary Three-particle GHZ State

We first introduce the process of realizing teleporaion of the three-particle unknown GHZ state. Suppose Alice is the sender and the two possible receivers are Bob and Charlie. If Bob is the receiver, then Charlie is the controller, and vice versa. The arbitrary GHZ state to be teleported is denoted by:

$$|\eta\rangle_{ABC} = (x|000\rangle + y|111\rangle)_{ABC} \quad (1)$$

where the unknown numbers  $x$  and  $y$  meet  $|x|^2 + |y|^2 = 1$ . A four-particle partially entangled state and a three-particle partially entangled state constitute the channel we need for teleportation. To ensure generality, the quantum channel linking Alice, Bob, and Charlie can be written as follows:

$$|\eta\rangle_{1234} = (a|0101\rangle + b|0110\rangle + c|1001\rangle + d|1010\rangle)_{1234} \quad (2)$$

$$|\eta\rangle_{567} = (m|011\rangle + n|100\rangle)_{567} \quad (3)$$

where  $a, b, c, d, m, n$  are real coefficients and satisfy  $|a|^2 + |b|^2 + |c|^2 + |d|^2 = 1$ ,  $|m|^2 + |n|^2 = 1$  and  $|a| > |b| > |c| > |d|$ ,  $|m| > |n|$ . The definition of Bell states  $|\Psi^\pm\rangle_{ij}$  and  $|\Phi^\pm\rangle_{ij}$  are given as following:

$$|\Psi^\pm\rangle_{ij} = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle)_{ij} \quad (4)$$

$$|\Phi^\pm\rangle_{ij} = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle)_{ij} \quad (5)$$

The above four-particle and three-particle partially entangled states are shared by two possible receivers (Bob and Charlie) and the sender (Alice) that are sufficiently far apart in space. Assume that the sender Alice has particles 1, 5, A, B, and C, Bob has particles 4,7, the remaining two particles 3 and 6, belong to Charlie. Initially, the state of the three-particle unknown GHZ state and the channel composed of four-particle and three-particle partially entangled state is given by:

$$|\eta\rangle = |\eta\rangle_{ABC} \otimes |\eta\rangle_{1234} \otimes |\eta\rangle_{567} \quad (6)$$

In our quantum teleportation scheme, there are

the following three steps.

1) In order to achieve teleportation, the first step Alice has to do is to carry out Bell-state measurement on particle pair (A,1) and particle pair (B,5) in turn. After the measurement is completed, the following different quantum states will be generated according to the different measurement basis:

$${}_{B5}\langle\Phi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{2}(xam|010111\rangle + xbm|011011\rangle \pm^2 \pm^1 ycn|100100\rangle \pm^2 \pm^1 ydn|101000\rangle)_{C23467} \quad (7)$$

$${}_{B5}\langle\Psi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{2}(xan|010100\rangle + xbn|011000\rangle \pm^2 \pm^1 ycm|100111\rangle \pm^2 \pm^1 ydm|101011\rangle)_{C23467} \quad (8)$$

$${}_{B5}\langle\Phi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{2}(xcm|000111\rangle + xdm|001011\rangle \pm^2 \pm^1 yan|110100\rangle \pm^2 \pm^1 ybn|111000\rangle)_{C23467} \quad (9)$$

$${}_{B5}\langle\Psi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{2}(xcn|000100\rangle + xdn|001000\rangle \pm^2 \pm^1 yam|110111\rangle \pm^2 \pm^1 ybm|111011\rangle)_{C23467} \quad (10)$$

where  $\pm^1, \pm^2$  represent the results corresponding to the measurement outcomes of the Bell-state on particle pairs (A,1) and (B,5), respectively.

2) Alice performs a single-particle measurement on particle C, where the measurement basis is  $|\xi^\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$ . After the measurement is completed, Eqs. (7)~Eqs. (10) can be rewritten as the following four equations:

$${}_C\langle\xi^\pm|_{B5}\langle\Phi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{2\sqrt{2}}(xam|10111\rangle + xbm|11011\rangle \pm^3 \pm^2 \pm^1 ycn|00100\rangle \pm^3 \pm^2 \pm^1 ydn|01000\rangle)_{23467} \quad (11)$$

$${}_C\langle\xi^\pm|_{B5}\langle\Psi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{2\sqrt{2}}(xan|10100\rangle + xbn|11000\rangle \pm^3 \pm^2 \pm^1 ycm|00111\rangle \pm^3 \pm^2 \pm^1 ydm|01011\rangle)_{23467} \quad (12)$$

$${}_C\langle\xi^\pm|_{B5}\langle\Phi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{2\sqrt{2}}(xcm|00111\rangle + xdm|01011\rangle \pm^3 \pm^2 \pm^1 yan|10100\rangle \pm^3 \pm^2 \pm^1 ybn|11000\rangle)_{23467} \quad (13)$$

$${}_C\langle\xi^\pm|_{B5}\langle\Psi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{2\sqrt{2}}(xcn|00100\rangle + xdn|01000\rangle \pm^3 \pm^2 \pm^1 yam|10111\rangle \pm^3 \pm^2 \pm^1 ybm|11011\rangle)_{23467} \quad (14)$$

where  $\pm^3$  denotes measurement result on particle C.

3) Now, notice that the attribution of particle 2 depends on the person who wants to restore the state  $|\eta\rangle_{ABC}$ , here we assume that Bob wants to restore the state  $|\eta\rangle_{ABC}$ , then Charlie is the controller. Alice has to transmit the measurements result to Bob and Charlie over the classical channel after she has completed the measurement, and Charlie needs to perform a Bell-state measurement on particles (3,6) at the request of Bob. After the measurement is completed, Eqs. (11)~Eqs. (14) can be rewritten as the following eight equations:

$${}_{36}\langle\Phi^\pm|_C\langle\xi^\pm|_{B5}\langle\Phi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{4}(\pm^4 xbm|101\rangle \pm^3 \pm^2 \pm^1 ycn|010\rangle)_{247} \quad (15)$$

$${}_{36}\langle\Psi^\pm|_C\langle\xi^\pm|_{B5}\langle\Phi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{4}(xam|111\rangle \pm^4 \pm^3 \pm^2 \pm^1 ydn|000\rangle)_{247} \quad (16)$$

$${}_{36}\langle\Phi^\pm|_C\langle\xi^\pm|_{B5}\langle\Psi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{4}(xan|110\rangle \pm^4 \pm^3 \pm^2 \pm^1 ydm|001\rangle)_{247} \quad (17)$$

$${}_{36}\langle\Psi^\pm|_C\langle\xi^\pm|_{B5}\langle\Psi^\pm|_{A1}\langle\Phi^\pm|\eta\rangle = \frac{1}{4}(\pm^4 xbn|100\rangle \pm^3 \pm^2 \pm^1 ycm|011\rangle)_{247} \quad (18)$$

$${}_{36}\langle\Phi^\pm|_C\langle\xi^\pm|_{B5}\langle\Phi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{4}(\pm^4 xdm|001\rangle \pm^3 \pm^2 \pm^1 yan|110\rangle)_{247} \quad (19)$$

$${}_{36}\langle\Psi^\pm|_C\langle\xi^\pm|_{B5}\langle\Phi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{4}(xcm|011\rangle \pm^4 \pm^3 \pm^2 \pm^1 ybn|100\rangle)_{247} \quad (20)$$

$${}_{36}\langle\Phi^\pm|_C\langle\xi^\pm|_{B5}\langle\Psi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{4}(xcn|010\rangle \pm^4 \pm^3 \pm^2 \pm^1 ybm|101\rangle)_{247} \quad (21)$$

$${}_{36}\langle\Psi^\pm|_C\langle\xi^\pm|_{B5}\langle\Psi^\pm|_{A1}\langle\Psi^\pm|\eta\rangle = \frac{1}{4}(\pm^4 xdn|000\rangle \pm^3 \pm^2 \pm^1 yam|111\rangle)_{247} \quad (22)$$

where  $\pm^4$  denotes Bell-state measurement outcomes corresponding to particles (3,6).

4) After the measurement, according to the different measurement results delivered by Charlie through the classical channel, Bob can choose the appropriate unitary operation to restore the unknown state  $|\eta\rangle_{ABC}$ . Then Bob needs to construct a corresponding relationship between  $|000\rangle_{247}, |111\rangle_{247}$

and the coefficients  $x$  and  $y$ , respectively. For instance, assume that the measurement results of Alice are  $|\Phi^+\rangle_{A1}|\Phi^+\rangle_{B5}|\xi^+\rangle_C$ , the measurement result of Charles is  $|\Phi^+\rangle_{36}$ , so when Charlie tells Bob the measurement result on particle pair (3,6), Bob can infer that the quantum state collapses into  $\frac{1}{4}(xbm|101\rangle + ycn|010\rangle)_{247}$ . To recover the initial state, Bob ought to implement the Pauli  $X$  operator (expressed as  $X$ ) on particle 5, 7 and perform the identity operator (expressed as  $I$ ) on particle 6. That is,  $U_1 = X_5 \otimes I_6 \otimes X_7$ . Therefore, Bob will obtain:

$$U_1_{(36)}\langle\Phi^+|_C\langle\xi^+|_{B5}\langle\Phi^+|_{A1}\langle\Phi^+|\eta\rangle = \frac{1}{4}(xbm|000\rangle + ycn|111\rangle)_{247} \quad (23)$$

5) Bob adds an auxiliary particle  $H$  where  $|0\rangle_H$  is as its original state. Then he performs an unitary matrix  $U_2$  based on a set of basis  $\{|00\rangle_{7H}, |01\rangle_{7H}, |10\rangle_{7H}, |11\rangle_{7H}\}$ :

$$U_2 = \begin{pmatrix} H_1 & H_2 \\ H_2 & -H_1 \end{pmatrix}$$

where  $H_i (i = 1, 2)$  is a  $2 \times 2$  matrix, they can be written as  $H_1 = \text{diag}(h_0, h_1)$ ,  $H_2 = \text{diag}(\sqrt{1-h_0^2}, \sqrt{1-h_1^2})$ , and  $(h_0, h_1) = (\frac{dn}{bm}, \frac{d}{c})$ . Under this transformation, the state in Eq.(23) will transform into:

$$\frac{1}{4}dn(x|000\rangle + y|111\rangle)_{247}|0\rangle_H + \frac{1}{4}[\sqrt{(bm)^2 - (dn)^2}x|000\rangle + n\sqrt{c^2 - d^2}y|111\rangle]_{247}|1\rangle_H \quad (24)$$

6) Finally, Bob needs to measure particle  $H$ . The measurement result of  $|0\rangle_H$  means that the teleportation of their joint efforts is successful and the measurement result of  $|1\rangle_H$  means failure.

In the process, Bob must carry out different unitary operations according to the state that he has got. Table 1 shows all possible states obtained by Bob and the corresponding unitary operation  $U_1$ . Table 2 shows the values of different  $h_i$  in unitary operation  $U_2$ .

The main process of our PQT scheme, in which Bob is as the receiver, is illustrated as Fig. 1. The particles are represented by solid circles and the entanglement between particles is represented by solid

lines. The drawings on the rightside are entanglement representation in steps 1), 2), 3), respectively. BSM and SM are as Bell-state measurement and single-particle measurement.

**Table 1 States and the corresponding unitary operation  $U_1$  for Bob**

Bob's states	Bob's operation
$xbm 101\rangle + ycn 010\rangle$	$X_5 \otimes I_6 \otimes X_7$
$xbm 101\rangle - ycn 010\rangle$	$X_5 \otimes I_6 \otimes (iY)_7$
$-xbm 101\rangle + ycn 010\rangle$	$(-iY)_5 \otimes I_6 \otimes X_7$
$-xbm 101\rangle - ycn 010\rangle$	$(-X)_5 \otimes I_6 \otimes X_7$
$xam 111\rangle + ydn 000\rangle$	$X_5 \otimes X_6 \otimes X_7$
$xam 111\rangle - ydn 000\rangle$	$X_5 \otimes X_6 \otimes (iY)_7$
$xan 110\rangle + ydm 001\rangle$	$X_5 \otimes X_6 \otimes I_7$
$xan 110\rangle - ydm 001\rangle$	$X_5 \otimes X_6 \otimes Z_7$
$xbn 100\rangle + ycm 011\rangle$	$X_5 \otimes I_6 \otimes I_7$
$xbn 100\rangle - ycm 011\rangle$	$X_5 \otimes I_6 \otimes Z_7$
$-xbn 100\rangle + ycm 011\rangle$	$(-iY)_5 \otimes I_6 \otimes I_7$
$-xbn 100\rangle - ycm 011\rangle$	$(-X)_5 \otimes I_6 \otimes I_7$
$xdm 001\rangle + yan 110\rangle$	$I_5 \otimes I_6 \otimes X_7$
$xdm 001\rangle - yan 110\rangle$	$I_5 \otimes I_6 \otimes (iY)_7$
$-xdm 001\rangle + yan 110\rangle$	$I_5 \otimes I_6 \otimes (-iY)_7$
$-xdm 001\rangle - yan 110\rangle$	$I_5 \otimes I_6 \otimes (-X)_7$
$xcm 011\rangle + ybn 100\rangle$	$I_5 \otimes X_6 \otimes X_7$
$xcm 011\rangle - ybn 100\rangle$	$I_5 \otimes X_6 \otimes (iY)_7$
$xcn 010\rangle + ybm 101\rangle$	$I_5 \otimes X_6 \otimes I_7$
$xcn 010\rangle - ybm 101\rangle$	$I_5 \otimes (iY)_6 \otimes I_7$
$xdn 000\rangle + yam 111\rangle$	$I_5 \otimes I_6 \otimes I_7$
$xdn 000\rangle - yam 111\rangle$	$I_5 \otimes I_6 \otimes Z_7$
$-xdn 000\rangle + yam 111\rangle$	$I_5 \otimes I_6 \otimes (-Z)_7$
$-xdn 000\rangle - yam 111\rangle$	$I_5 \otimes I_6 \otimes (-I)_7$

**Table 2  $h_i$  in the unitary operation  $U_2$**

restored state by $U_1$	$h_0$	$h_1$
$xbm 000\rangle + ycn 111\rangle$	$\frac{dn}{bm}$	$\frac{d}{c}$
$xam 000\rangle + ydn 111\rangle$	$\frac{dn}{am}$	1
$xan 000\rangle + ydm 111\rangle$	$\frac{d}{a}$	$\frac{n}{m}$
$xbn 000\rangle + ycm 111\rangle$	$\frac{d}{b}$	$\frac{dn}{cm}$
$xdm 000\rangle + yan 111\rangle$	$\frac{n}{m}$	$\frac{d}{a}$
$xcm 000\rangle + ybn 111\rangle$	$\frac{dn}{cm}$	$\frac{d}{b}$
$xcn 000\rangle + ybm 111\rangle$	$\frac{d}{c}$	$\frac{dn}{bm}$
$xdn 000\rangle + yam 111\rangle$	1	$\frac{dn}{am}$




Fig. 1 A flow chart of our PQT scheme

In the former case, the state of particles 2,4,7 will collapse into  $\frac{1}{4}dn(x|000) + y|111\rangle_{247}$ , and the probability of success is  $\frac{1}{16}d^2n^2$ . Simultaneously, for Eq. (15), since Alice and Charlie have two different choices for each measurement, there are 16 possible cases in total. It's easy to figure out that the probability of success is  $d^2n^2$ . Considering that there are 128 cases, the total success probability is  $8d^2n^2$ . We show the success probability surface of our PQT in Fig. 2, which gives the relationship between probability and coefficients  $d$  and  $n$ .




Fig. 2 Success probability surface of our PQT

If the coefficients in Eq. (2) satisfy  $a = b = c = d = \frac{1}{2}$  and the coefficients in Eq. (3) satisfy  $m = n = \frac{1}{\sqrt{2}}$ , then the channel will be in its maximally entangled states, and thus the probability of teleportation equals 1. Namely, this is the usual teleportation. Fig. 3 illustrates the relation between  $d$

and  $n$  when the probability  $8d^2n^2 = 1$ .




Fig. 3 Probability  $8d^2n^2 = 1$

Likewise, if the person who wants to restore the state  $|\eta\rangle_{ABC}$  is Charlie, then assign particle 2 to him. First of all, after Alice makes her measurements on particles  $(A,1)$ ,  $(B,5)$ , and  $C$ , she sends the measurement result to Charlie through the classical channel. Next, Bob needs to carry out a Bell-state measurement on particles  $(4,7)$  at the request of Charlie. The remaining steps are the same as the case when the person who wants to restore the state  $|\eta\rangle_{ABC}$  is Bob. It is not difficult to figure out the probability that Charlie can recover the initial state  $|\eta\rangle_{ABC}$  is  $8d^2n^2$ . Accordingly, we can find that whether the person who wants to restore the original state is Bob or Charlie, the probability of success is the same in both cases.

## 2 Probabilistic Teleportation of Arbitrary Three-particle W State

Assuming that the state we want to transmit is not the GHZ state but the three-particle W state, through the following discussion we can find that the quantum channel we constructed in Eqs. (2) and Eqs. (3) can also realize the teleportation of arbitrary three-particle W states. The arbitrary W state to be teleported can be written as:

$$|\eta'\rangle_{A'B'C'} = (p|001\rangle + q|010\rangle + r|100\rangle)_{A'B'C'} \quad (25)$$

where the unknown real numbers  $p, q, r$  satisfy  $|p|^2 + |q|^2 + |r|^2 = 1$ .

If Alice is the sender, she intends to transmit a three-particle W state, and Charlie, who is far away from Alice, wants to get this state with the help of Bob (Here, for simplicity, we only consider the cases where Charlie is the receiver). Assume that Alice has particles 1,5, A, B, and C, Charlie has particles 2,4,7

and the remaining two particles 3,6 belong to the controller Bob. Initially, the state of the composite system is given by:

$$|\eta'\rangle = |\eta'\rangle_{A'B'C'} \otimes |\eta\rangle_{1234} \otimes |\eta\rangle_{567} \quad (26)$$

To achieve teleportation from Alice to Charlie, the first step Alice needs to do is to operate Bell-state measurement on particle pairs (A',1) and (B',5) in possession, respectively. Next, Alice needs to perform single-particle measurement on particle C, where the measurement basis is  $|\xi^{\pm}\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$ . After

operations, Alice needs to send the measurement result to Bob and Charlie through the classical channel. If Bob is willing to help Charlie to rebuild the initial state, Bob has to carry out Bell-state measurement on particle pair (3,6). In the light of the result of Alice and Bob, Charlie will be able to rebuild the original state in Eq. (25) by performing a suitable unitary operation  $U'_1$  on the particles 2,4,7. We show all possible states obtained by Charlie and the corresponding unitary operation  $U'_1$  in Table 3.

**Table 3** Unitary operations  $U'_1$  corresponding to the state obtained by Charlie

Charlie's states	Charlie's operation
$\pm^4 \pm^3 pbm 101\rangle \pm^2 qan 110\rangle \pm^4 \pm^1 rdm 001\rangle$	$ 000\rangle\langle 000  +  110\rangle\langle 010  +  101\rangle\langle 100  \pm^2  010\rangle\langle 110  \pm^4 \pm^1  100\rangle\langle 001  +  011\rangle\langle 011  \pm^4 \pm^3  001\rangle\langle 101  +  111\rangle\langle 111 $
$\pm^3 pam 111\rangle \pm^4 \pm^2 qbn 100\rangle \pm^1 rcm 011\rangle$	$ 000\rangle\langle 000  +  011\rangle\langle 010  \pm^4 \pm^2  010\rangle\langle 100  +  110\rangle\langle 110  +  101\rangle\langle 001  \pm^1  100\rangle\langle 011  +  111\rangle\langle 101  \pm^3  001\rangle\langle 111 $
$\pm^3 pan 110\rangle \pm^4 \pm^2 qbm 101\rangle \pm^1 rcn 010\rangle$	$ 000\rangle\langle 000  \pm^1  100\rangle\langle 010  +  110\rangle\langle 100  \pm^3  001\rangle\langle 110  +  101\rangle\langle 001  +  011\rangle\langle 011  \pm^4 \pm^2  010\rangle\langle 101  +  111\rangle\langle 111 $
$\pm^4 \pm^3 pbn 100\rangle \pm^2 qam 111\rangle \pm^4 \pm^1 rdn 000\rangle$	$ 000\rangle\langle 001  +  011\rangle\langle 011  +  101\rangle\langle 101  \pm^2  010\rangle\langle 111  \pm^4 \pm^1  100\rangle\langle 000  +  110\rangle\langle 010  \pm^4 \pm^3  001\rangle\langle 100  +  111\rangle\langle 110 $
$\pm^4 \pm^3 pdm 001\rangle \pm^2 qcn 010\rangle \pm^4 \pm^1 rbm 101\rangle$	$ 000\rangle\langle 000  \pm^2  010\rangle\langle 010  +  101\rangle\langle 100  +  110\rangle\langle 110  \pm^4 \pm^3  001\rangle\langle 001  +  011\rangle\langle 011  \pm^4 \pm^1  100\rangle\langle 101  +  111\rangle\langle 111 $
$\pm^3 pcm 011\rangle \pm^4 \pm^2 qdn 000\rangle \pm^1 ram 111\rangle$	$ 001\rangle\langle 010  \pm^4 \pm^2  010\rangle\langle 000  +  000\rangle\langle 100  +  110\rangle\langle 110  +  111\rangle\langle 001  \pm^3  011\rangle\langle 011  +  101\rangle\langle 101  \pm^1  100\rangle\langle 111 $
$\pm^3 pcn 010\rangle \pm^4 \pm^2 qdm 001\rangle \pm^1 ran 110\rangle$	$ 011\rangle\langle 011  \pm^4 \pm^2  010\rangle\langle 001  +  101\rangle\langle 101  +  111\rangle\langle 111  +  000\rangle\langle 000  \pm^3  001\rangle\langle 010  +  110\rangle\langle 100  \pm^1  100\rangle\langle 110 $
$\pm^4 \pm^3 pdn 000\rangle \pm^2 qcm 011\rangle \pm^4 \pm^1 rbn 100\rangle$	$ 000\rangle\langle 001  \pm^2  010\rangle\langle 011  +  101\rangle\langle 101  +  111\rangle\langle 111  \pm^4 \pm^3  001\rangle\langle 000  +  011\rangle\langle 010  \pm^4 \pm^1  100\rangle\langle 100  +  110\rangle\langle 110 $

Similarly, Charlie needs to add an auxiliary particle  $H$ ,  $|0\rangle_H$  is its original state. Then Charlie performs unitary matrix  $U'_2$  based on a set of basis  $\{|000\rangle_{47H}, |100\rangle_{47H}, |010\rangle_{47H}, |001\rangle_{47H}, |101\rangle_{47H}, |011\rangle_{47H}\}$ :

$$U'_2 = \begin{pmatrix} H'_1 & H'_2 \\ H'_2 & -H'_1 \end{pmatrix}$$

where  $H'_i$  ( $i = 1,2$ ) is a  $3 \times 3$  matrix, written as  $H'_1 = \text{diag}(h_0, h_1, h_2)$ ,  $H'_2 = \text{diag}(\sqrt{1-h_0^2}, \sqrt{1-h_1^2}, \sqrt{1-h_2^2})$ . The values of  $h_i$  in the unitary operations  $U'_2$  are shown in Table 4.

Since the measurement process is the same as Section 1, we only give the collapse state of particles (2,4,7) after measurement.

**Table 4**  $h_i$  in the unitary operations  $U'_2$

restored state by $U'_1$	$h_0$	$h_1$	$h_2$
$pbm 001\rangle + qan 010\rangle + rdm 100\rangle$	$\frac{n}{m}$	$\frac{d}{a}$	$\frac{dn}{bm}$
$pam 001\rangle + qbn 010\rangle + rcm 100\rangle$	$\frac{dn}{cm}$	$\frac{d}{b}$	$\frac{dn}{am}$
$pan 001\rangle + qbm 010\rangle + rcn 100\rangle$	$\frac{d}{c}$	$\frac{dn}{bm}$	$\frac{d}{a}$
$pbn 001\rangle + qam 010\rangle + rdn 100\rangle$	1	$\frac{dn}{am}$	$\frac{d}{b}$
$pdm 001\rangle + qcn 010\rangle + rbm 100\rangle$	$\frac{dn}{bm}$	$\frac{d}{c}$	$\frac{n}{m}$
$pcm 001\rangle + qdn 010\rangle + ram 100\rangle$	$\frac{dn}{am}$	1	$\frac{dn}{cm}$
$pcn 001\rangle + qdm 010\rangle + ran 100\rangle$	$\frac{d}{a}$	$\frac{n}{m}$	$\frac{d}{c}$
$pdn 001\rangle + qcm 010\rangle + rbn 100\rangle$	$\frac{d}{b}$	$\frac{dn}{cm}$	1

$$\frac{1}{4}(\pm^4 \pm^3 p b m |101\rangle \pm^2 q a n |110\rangle \pm^4 \pm^1 r d m |001\rangle)_{247} \quad (27)$$

$$\frac{1}{4}(\pm^3 p a m |111\rangle \pm^4 \pm^2 q b n |100\rangle \pm^1 r c m |011\rangle)_{247} \quad (28)$$

$$\frac{1}{4}(\pm^3 p a n |110\rangle \pm^4 \pm^2 q b m |101\rangle \pm^1 r c n |010\rangle)_{247} \quad (29)$$

$$\frac{1}{4}(\pm^4 \pm^3 p b n |100\rangle \pm^2 q a m |111\rangle \pm^4 \pm^1 r d n |000\rangle)_{247} \quad (30)$$

$$\frac{1}{4}(\pm^4 \pm^3 p d m |001\rangle \pm^2 q c n |010\rangle \pm^4 \pm^1 r b m |101\rangle)_{247} \quad (31)$$

$$\frac{1}{4}(\pm^3 p c m |011\rangle \pm^4 \pm^2 q d n |000\rangle \pm^1 r a m |111\rangle)_{247} \quad (32)$$

$$\frac{1}{4}(\pm^3 p c n |010\rangle \pm^4 \pm^2 q d m |001\rangle \pm^1 r a n |110\rangle)_{247} \quad (33)$$

$$\frac{1}{4}(\pm^4 \pm^3 p d n |000\rangle \pm^2 q c m |011\rangle \pm^4 \pm^1 r b n |100\rangle)_{247} \quad (34)$$

Now we take  $|\Phi^+\rangle_{A1}|\Phi^+\rangle_{B5}|\xi^+\rangle_C|\Phi^+\rangle_{36}$  as an example to show how to achieve teleportation. Suppose that:

$$\frac{1}{4}(p b m |101\rangle + q a n |110\rangle + r d m |001\rangle)_{247} \quad (35)$$

In order to recover the initial state  $|\eta'\rangle_{A'B'C}$ , Charlie has to carry out an appropriate unitary operation:

$$U'_1 = |000\rangle\langle 000| + |110\rangle\langle 010| + |101\rangle\langle 100| + |010\rangle\langle 110| + |100\rangle\langle 001| + |011\rangle\langle 011| + |001\rangle\langle 101| + |111\rangle\langle 111| \quad (36)$$

Then, Charlie will obtain:

$$\frac{1}{4}(p b m |001\rangle + q a n |010\rangle + r d m |100\rangle)_{247} \quad (37)$$

Next, we set  $h_0 = \frac{n}{m}$ ,  $h_1 = \frac{d}{a}$ ,  $h_2 = \frac{d n}{b m}$  in the unitary matrix  $U'_2$ . Through a unitary transformation similar to  $U_2$ , the state in Eq. (27) will transform into:

$$\frac{1}{4} d n (p |001\rangle + q |010\rangle + r |100\rangle)_{247} |0\rangle_H + \frac{1}{4} [\sqrt{(b m)^2 - (c n)^2} p |001\rangle + n \sqrt{a^2 - b^2} q |010\rangle + d \sqrt{m^2 - n^2} r |100\rangle]_{247} |1\rangle_H \quad (38)$$

Finally, Charlie needs to measure particle  $H$ . The measurement result of  $|0\rangle_H$  means the teleportation of their joint efforts is successful, and the measurement result of  $|1\rangle_H$  means failure. In the former case, the state of particles 2, 4 and 7 will collapse into  $\frac{1}{4} d n (p |001\rangle + q |010\rangle + r |100\rangle)_{247}$ , and the probability of success is  $\frac{1}{16} d^2 n^2$ . Similarly, other possible states can be calculated in the same way. Therefore, the total probability of success is  $8 d^2 n^2$  based on all 128 cases. If the coefficients in Eq. (2) satisfy  $a = b = c = d = \frac{1}{2}$  and the coefficients in Eq. (3) satisfy  $m = n = \frac{1}{\sqrt{2}}$ , then the channel will be in its maximally entangled state, and the probability of teleportation equals 1.

### 3 Conclusions

In summary, two explicit tripartite schemes are proposed for the controllable probabilistic teleportation of any unknown partially entangled GHZ or W state. At the same time, our schemes can implement teleportation from the sender to either one of the two receivers. In these two schemes, we construct a new quantum channel which is consisted of a one-dimensional four-particle partially entangled state and a three-particle partially entangled state. This channel can realize not only the transmission of the GHZ state but also the transmission of the W state. The processes of our schemes are the sender carries out two Bell-state measurement and a single-particle measurement on suitable particles, the controller performs a Bell-state measurement, the receiver performs the corresponding unitary operation and introduces an auxiliary particle. After these processes are completed, the unknown state can be transmitted with a certain probability, depending on the coefficient with the smallest absolute value in the four-particle partially entangled state and the coefficient with the smallest absolute value in the three-particle partially entangled state.

Since devices cannot be perfect in the experiment, the generation of maximally entangled state is not

possible. Entanglement purification methods are usually helpful for acquiring the ideal entangled state but often requires additional resources. Accordingly, our schemes are more feasible in experiments. We can find that the overall success probabilities of the two schemes are both  $8d^2n^2$ . If the channel is the maximally entangled state, then probabilistic teleportation becomes the usual teleportation, which has a probability of success equal to 100%.

### References

- [1] EKERT A K. Quantum cryptography based on Bell's theorem[J]. *Physical Review Letters*, 1991, 67(6): 661.
- [2] LIU B, XIA S, XIAO D, et al. Decoy-state method for quantum-key-distribution-based quantum private query[J]. *Science China Physics, Mechanics & Astronomy*, 2022, 65(4): 240312.
- [3] LI Z J, WEI K J. Improving parameter optimization in decoy-state quantum key distribution[J]. *Quantum Engineering*, 2022, 2022: 1-9.
- [4] BENNETT C H, BRASSARD G, POPESCU S, et al. Purification of noisy entanglement and faithful teleportation via noisy channels[J]. *Physical Review Letters*, 1996, 76(5): 722.
- [5] DENG F G, LONG G L, LIU X S. Two-step quantum direct communication protocol using the Einstein-Podolsky-Rosen pair block[J]. *Physical Review A*, 2003, 68(4): 042317.
- [6] JOY D, SURENDRAN S P, SABIR M. Efficient deterministic secure quantum communication protocols using multipartite entangled states[J]. *Quantum Information Processing*, 2017, 16: 1-11.
- [7] ZHOU L, SHENG Y B. One-step device-independent quantum secure direct communication[J]. *Science China Physics, Mechanics & Astronomy*, 2022, 65(5): 250311.
- [8] BENNETT C H, BRASSARD G, CRÉPEAU C, et al. Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels[J]. *Physical Review Letters*, 1993, 70(13): 1895.
- [9] IKRAM M, ZHU S Y, ZUBAIRY M S. Quantum teleportation of an entangled state[J]. *Physical Review A*, 2000, 62(2): 022307.
- [10] DI T, MUTHUKRISHNAN A, SCULLY M O, et al. Quantum teleportation of an arbitrary superposition of atomic Dicke states[J]. *Physical Review A*, 2005, 71(6): 062308.
- [11] PIRANDOLA S, EISERT J, WEEDBROOK C, et al. Advances in quantum teleportation[J]. *Nature Photonics*, 2015, 9(10): 641-652.
- [12] CHEN X, SHEN Y, ZHANG F L. Perfect teleportation with a partially entangled quantum channel[J]. *Physical Review A*, 2022, 106(3): 032430.
- [13] VAIDMAN L. Teleportation of quantum states[J]. *Physical Review A*, 1994, 49(2): 1473.
- [14] BOUWMEESTER D, PAN J W, MATTLE K, et al. Experimental quantum teleportation[J]. *Nature*, 1997, 390(6660): 575-579.
- [15] BRAUNSTEIN S L, KIMBLE H J. Teleportation of continuous quantum variables[J]. *Physical Review Letters*, 1998, 80(4): 869.
- [16] LEE H W, KIM J. Quantum teleportation and Bell's inequality using single-particle entanglement[J]. *Physical Review A*, 2000, 63(1): 012305.
- [17] YUAN H C, QI K G. Quantum logic networks for controlled teleportation of a single particle via W state[J]. *Chinese Physics*, 2005, 14(5): 898.
- [18] TAN X Q, YANG P, SONG T T. Teleportation of three-particle W state[J]. *International Journal of Theoretical Physics*, 2018, 57: 329-338.
- [19] LI D F, WANG R J, BAAGYERE E. Quantum teleportation of an arbitrary two-qubit state by using two three-qubit GHZ states and the six-qubit entangled state[J]. *Quantum Information Processing*, 2019, 18: 1-15.
- [20] ZHENG T, CHANG Y, ZHANG S B. Arbitrated quantum signature scheme with quantum teleportation by using two three-qubit GHZ states[J]. *Quantum Information Processing*, 2020, 19: 1-15.
- [21] AGRAWAL P, PATI A K. Probabilistic quantum teleportation[J]. *Physics Letters A*, 2002, 305(1-2): 12-17.
- [22] PATI A K, AGRAWAL P. Probabilistic teleportation and quantum operation[J]. *Journal of Optics B: Quantum and Semiclassical Optics*, 2004, 6(8): S844.
- [23] GAO T, YAN F L, LI Y C. Optimal controlled teleportation[J]. *Europhysics Letters*, 2008, 84(5): 50001.
- [24] GORDON G, RIGOLIN G. Generalized quantum-state sharing[J]. *Physical Review A*, 2006, 73(6): 062316.
- [25] LUO Q M, ZHANG T G, HUANG X F, et al. Two quantum proxy blind signature schemes based on controlled quantum teleportation[J]. *Entropy*, 2022, 24(10): 1421.
- [26] WANG T Y, WEN Q Y. Controlled quantum teleportation with Bell states[J]. *Chinese Physics B*, 2011, 20(4): 040307.
- [27] WANG M T, LI H S. Bidirectional quantum teleportation using a five-qubit cluster state as a quantum channel[J]. *Quantum Information Processing*, 2022, 21(2): 44.
- [28] THAPLIYAL K, VERMA A, PATHAK A. A general method for selecting quantum channel for bidirectional controlled state teleportation and other schemes of controlled quantum communication[J]. *Quantum Information Processing*, 2015, 14: 4601-4614.
- [29] DAI R, LI H S. Asymmetric bidirectional quantum teleportation via seven-qubit cluster state[J]. *International Journal of Theoretical Physics*, 2022, 61(7): 187.
- [30] VERMA V, YADAV D, MISHRA D K. Improvement on cyclic controlled teleportation by using a seven-qubit entangled state[J]. *Optical and Quantum Electronics*, 2021, 53: 1-11.
- [31] VERMA V. Symmetric and asymmetric cyclic controlled quantum teleportation via nine-qubit entangled state[J]. *Modern Physics Letters B*, 2021, 35(15): 2150249.
- [32] YANG B C. Asymmetric cyclic controlled quantum teleportation by using a quantum channel composed of six G states[J]. *Modern Physics Letters B*, 2022, 36(22): 2250113.